arXiv:1211.0687v1 [math.AG] 4 Nov 2012

MIXED HODGE STRUCTURES AND WEIERSTRASS σ-FUNCTION

GRZEGORZ BANASZAK* AND JAN MILEWSKI

Abstract. Un σ-opérateur sur la complexification $V_{\mathbb{C}}$ d'un espace vectoriel réel $V_{\mathbb{R}}$ est un opérateur $A \in End_{\mathbb{C}}(V_{\mathbb{C}})$ tel que $\sigma(A) = 0$, où $\sigma(z)$ est la fonction σ de Weierstrass. Dans cet article, nous introduisons la notion de σ-opérateur fortement pseudo-réel et démontrons qu'il y a une correspondance biunivoque entre les structures de Hodge mixtes réelles et les σ-opérateurs fortement pseudo-réels.

A σ-operator on a complexification $V_{\mathbb{C}}$ of an $\mathbb{R}$-vector space $V_{\mathbb{R}}$ is an operator $A \in \mathrm{End}_{\mathbb{C}}(\mathrm{V}_{\mathbb{C}})$ such that $\sigma(A) = 0$ where $\sigma(z)$ denotes the Weierstrass σ-function. In this paper we define the notion of the strongly pseudo-real σ-operator and prove that there is one to one correspondence between real mixed Hodge structures and strongly pseudo-real σ-operators.

1. Introduction

Let $\mathbb{S} := R_{\mathbb{C}/\mathbb{R}}\mathbb{G}_m$ and let $V_{\mathbb{R}}$ be a finite dimensional real vector space. By a real Hodge structure (HS) on $V_{\mathbb{R}}$ we understand a finite direct sum of real pure Hodge structures with given weights cf. [1]. We can consider a Hodge structure as a real algebraic group representation $\rho : \mathbb{S} \longrightarrow \mathrm{GL}(V_{\mathbb{R}})$ (see eg. [1], [5], [6] for the definition of HS). Let $\mathcal{L}\rho$ be the Lie algebra representation of ρ. The following operator

$$S := S(\rho) := \mathcal{L}\rho(1 + i) \in \mathrm{End}(V_{\mathbb{R}}),$$

introduced in [1], will be called the Hodge-Lie operator of the real HS given by ρ.

Let $\sigma(z)$ be the Weierstrass' sigma function for the lattice $\Lambda := \mathbb{Z}\,\omega_1 + \mathbb{Z}\,\omega_2$ with $\omega_1 = 1 - i$ and $\omega_2 = 1 + i$:

$$\sigma(z) := z \prod_{(p,q)\neq(0,0)} \left(1 - \frac{z}{\lambda_{p,q}}\right) \exp\left[\frac{z}{\lambda_{p,q}} + \frac{1}{2}\left(\frac{z}{\lambda_{p,q}}\right)^2\right]$$

where

$$\lambda_{p,q} := p\omega_1 + q\omega_2.$$

Definition 1.1. Let V_K ($K = \mathbb{R}$ resp. $K = \mathbb{C}$) be a finite dimensional vector space over K. An endomorphism $A \in \mathrm{End}(V_K)$ is called a (real resp. complex) σ-operator if $\sigma(A) = 0$.

Observe that if $V_{\mathbb{C}} = V_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$ then A is a σ-operator on $V_{\mathbb{C}}$ if and only if $\overline{A}$ is a σ-operator. In [1] we obtained the following result:

2010 *Mathematics Subject Classification.* 14D07.

Key words and phrases. Mixed Hodge structure, Deligne splitting, Weierstrass σ function.

*Partially supported by the NCN (National Center of Science for Poland) NN201 607440.

Theorem 1.2. (i) *Hodge-Lie operator S of any HS is a real σ-operator.*
(ii) *There is one to one correspondence between HS and the real σ-operators. This equivalence is given by assigning to a HS its Hodge-Lie operator:*

$$\rho \longleftrightarrow S(\rho)$$

Christopher Deninger asked us very interesting question whether our approach to real HS via Weierstrass σ function can be extended to the mixed Hodge structures (MHS). This paper is an affirmative answer to the C. Deninger's question (see Definition 1.5 and Theorem 1.6 below). For the definition of MHS see eg. [2], [3], [4] cf. the Definition 2.1 below. In all of this paper a MHS means a real MHS.

Let us introduce basic definitions and state the main results of this paper. The proofs of these results are given in section 3. For every $\lambda \in \Lambda$ define:

$$\sigma_{\lambda}(z) := \frac{\sigma(z)}{z-\lambda}. \tag{1}$$

We will write $\sigma_{m,n}(z) := \sigma_{\lambda_{m,n}}(z)$ for all $(m,n) \in \mathbb{Z} \times \mathbb{Z}$.

Definition 1.3. Let $V_{\mathbb{C}} := V_{\mathbb{R}} \otimes \mathbb{C}$. A σ-operator A on $V_{\mathbb{C}}$ is called a weakly pseudo-real if

$$\sigma_{r,s}(A)(\overline{A}-A)\sigma_{p,q}(A) = 0 \tag{2}$$

for all $r+s \geq p+q$.

Theorem 1.4. *Every weakly pseudo-real σ-operator A determines a MHS.*

A natural question is whether there is one to one correspondence between weakly pseudo-real σ-operators and real MHS on $V_{\mathbb{R}}$. In general the answer is no. However, if we strengthen the condition (2), then we can strengthen the Theorem 1.4.

Definition 1.5. Let $V_{\mathbb{C}} := V_{\mathbb{R}} \otimes \mathbb{C}$. A σ-operator A on $V_{\mathbb{C}}$ is called a strongly pseudo-real if

$$\sigma_{r,s}(A)(\overline{A}-A)\sigma_{p,q}(A) = 0 \tag{3}$$

for all $r \geq p$ or $s \geq q$.

Theorem 1.6. *There is one to one correspondence between MHS and strongly pseudo-real σ-operators.*

Observe that A is a weakly pseudo-real (resp. strongly pseudo-real) σ-operator if only if $\overline{A}$ is a weakly pseudo-real (resp. strongly pseudo-real) σ-operator.

2. Deligne splitting of a mixed Hodge structure

Definition 2.1. A real MHS on $V_{\mathbb{R}}$ consists of two filtrations, a finite increasing filtration on $V_{\mathbb{R}}$, the weight filtration ${W_{\bullet}}^{\mathbb{R}}$ and a finite decreasing filtration $F^{\bullet}$ on $V_{\mathbb{C}} := V_{\mathbb{R}} \otimes \mathbb{C}$, the Hodge filtration which induce a pure real HS of the weight n on each graded piece

$$\mathrm{Gr}_n^W(V_{\mathbb{R}}) = W_n^{\mathbb{R}}/W_{n-1}^{\mathbb{R}}. \tag{4}$$

Theorem 2.2. (Deligne splitting of MHS). *For any MHS there exists exactly one decomposition:*

$$V_{\mathbb{C}} = \bigoplus_{p,q} I^{p,q} \tag{5}$$

such that

$$W_n = \bigoplus_{p+q\leq n} I^{p,q}, \quad F^p = \bigoplus_{\substack{k\geq p\\ q\in\mathbb{Z}}} I^{k,q}, \tag{6}$$

$$I^{p,q} = \overline{I^{q,p}} \mod D_{p-1,q-1}, \tag{7}$$

where

$$W_n := W_n^{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}, \quad D_{r,s} := \bigoplus_{\substack{k\leq r\\ l\leq s}} I^{k,l}. \tag{8}$$

This decomposition can be expressed via weight and Hodge filtrations in the following way:

$$I^{p,q} = V^p_{p+q} \cap (\overline{V^q_{p+q}} + \overline{U^{q-1}_{p+q-1}}), \tag{9}$$

where

$$V^p_n := F^p \cap W_n, \qquad U^m_n := \sum_{j\geq 0} V^{m-j}_{n-j}. \tag{10}$$

For the proof of the Theorem 2.2 see [4] (cf. [2] pp. 471-472). Observe that $\overline{V^q_n} = \overline{F^q} \cap W_n$ because $\overline{W_n} = W_n$. Moreover

$$D_{p,q} = \overline{D_{q,p}} \tag{11}$$

$$W_n = \sum_{p+q=n} D_{p,q} \tag{12}$$

3. Mixed Hodge Structures via pseudo-real σ-operators

Let A be a σ-operator on $V_C = V_R \otimes \mathbb{C}$. We get the following decomposition into eigenspaces:

$$V_{\mathbb{C}} := \bigoplus_{p,q} I^{p,q}_A, \qquad I^{p,q}_A := \{x \in V : Ax = \lambda_{p,q} x\}. \tag{13}$$

Certainly

$$I\frac{p,q}{A} = \overline{I^{q,p}_A} \tag{14}$$

because $\lambda_{q,p} = \overline{\lambda}_{p,q}$. Define the weight and Hodge filtrations and bifiltration of this σ-operator in the following way:

$$W^A_n := \bigoplus_{p+q\leq n} I^{p,q}_A, \quad F^p_A := \bigoplus_{\substack{k\geq p\\ q\in\mathbb{Z}}} I^{k,q}_A, \quad D^A_{p,q} := \bigoplus_{\substack{k\leq p\\ l\leq q}} I^{k,l}_A. \tag{15}$$

Definition 3.1. Two σ-operators A_1 and A_2 are called weakly equivalent if they determine the same weight filtration and they induce the same homomorphism on the grading of the weight filtration (hence their difference is a homomorphism of weight filtration of degree -1).

Lemma 3.2. *Let $V_{\mathbb{C}}$ be a complexification of the real vector space $V_{\mathbb{R}}$. A σ-operator $A \in \mathrm{End}_{\mathbb{C}}(V_{\mathbb{C}})$ is weakly pseudo-real if and only if operators A and $\overline{A}$ are weakly equivalent.*

Proof. Consider the decomposition (13). The lemma is a consequence of the following equality:

$$\sigma_{p,q}(A) = \sigma_{p,q}(\lambda_{p,q})P_{p,q},$$

where $P_{p,q}$ denotes the projection operator

$$P_{p,q} : \bigoplus_{m,n} I_A^{m,n} \to I_A^{p,q}$$

onto the direct summand $I_A^{p,q}$. □

Proof of the Theorem 1.4.
By Lemma 3.2 the σ-operators A and $\overline{A}$ are weakly equivalent, hence they determine the same weight filtration $W_\bullet = \overline{W_\bullet}$, where $W_\bullet := W_\bullet^A$ and $\overline{W_\bullet} = W_\bullet^{\overline{A}}$. Hence the weight filtration $W_\bullet$ is a complexification of certain increasing filtration $W_\bullet^{\mathbb{R}}$ on $V_{\mathbb{R}}$. Induced quotient operators $[A]_n$, $[\overline{A}]_n$ on W_n/W_{n-1} are equal, hence are real σ-operators (a real σ-operator on $V_{\mathbb{C}}$ is a σ-operator which is a complexification of an operator on $V_{\mathbb{R}}$). All eigenvalues of the operator $[A]_n$ have real part equal n, hence this operator gives a pure Hodge structure of weight n. Hodge filtration of the σ-operator $[A]_n$ on W_n/W_{n-1} is induced by the Hodge filtration of A. Hence the weight and Hodge filtrations of the σ-operator A induce a mixed Hodge structure. □

The double increasing, finite filtration $D^A_{\bullet,\bullet}$ given in (15) leads to the following definition:

Definition 3.3. Two σ-operators A_1 and A_2 are called strongly equivalent if they determine the same double filtrations $D^{A_1}_{\bullet,\bullet} = D^{A_2}_{\bullet,\bullet}$ and their difference is a homomorphism of this bifiltration of bidegree $(-1,-1)$.

Lemma 3.4. *A σ-operator A is strongly pseudo-real if and only if operators A and $\overline{A}$ are strongly equivalent.*

Proof. The proof is very similar to the proof of Lemma 3.2 concerning the weakly pseudo-real σ-operator. □

Proof of the Theorem 1.6.
Every mixed Hodge structure $(W_\bullet, F^\bullet)$ determines uniquely the canonical Deligne's decomposition (5) - (8). Hence σ-operator A

$$A := \bigoplus_{p,q} \lambda_{p,q}\, \mathrm{id}_{I^{p,q}}$$

is also uniquely determined by this mixed Hodge structure and is strongly pseudo-real. Indeed

$$\overline{A} = \bigoplus_{p,q} \lambda_{p,q}\, \mathrm{id}_{\overline{I^{q,p}}},$$

because $\overline{\lambda}_{q,p} = \lambda_{p,q}$. Recall that $D_{q,p} = \overline{D}_{p,q}$. Let $x \in D_{p,q}$. For $r \le p,\ s \le q,\ k < r,\ l < s$ there exist vectors $\widetilde{x}_{r,s} \in \overline{I^{s,r}}$, $x_{r,s} \in I^{r,s}$, $u_{r,s;k,l} \in I^{k,l}$ such that

$$x = \sum_{\substack{r \le p \\ s \le q}} \widetilde{x}_{r,s}\ , \quad \widetilde{x}_{r,s} = x_{r,s} + \sum_{\substack{k < r \\ l < s}} u_{r,s;k,l}.$$

Hence

$$(\overline{A} - A)x = \sum_{r,s,k,l} (\lambda_{r,s} - \lambda_{k,l}) u_{r,s;k,l} \in D_{p-1,q-1},$$

where the sum $\sum_{r,s,k,l}$ is given for indices r, s, k, l in the range $r \le p,\ s \le q,\ k < r,\ l < s$. By Lemma 3.4 the σ-operator A is strongly pseudo-real.

Conversely every σ-operator A determines unique decomposition of $V_{\mathbb{C}}$ into eigensubspaces (see (13) and (15)). When σ-operator A is strongly pseudo-real then, by Lemma 3.4, equalities (13) and (15) give the Deligne canonical decomposition of the mixed Hodge structure determined by A. Indeed in this case for $x \in I_A^{p,q}$ we have $\overline{A}x - Ax =: u \in D^A_{p-1,q-1}$. Hence

$$\overline{A}x = \lambda_{p,q} x + u. \tag{16}$$

It remains to check that:

$$x + u' \in \overline{I_A^{q,p}}, \tag{17}$$

for some $u' \in D^A_{p-1,q-1}$. Since A is strongly pseudo-real we get $D^A_{\bullet,\bullet} = D^{\overline{A}}_{\bullet,\bullet}$. Observe that the operator $\lambda_{p,q}\mathrm{id} - \overline{A}$ with domain and target restricted to the invariant subspace $D^A_{p-1,q-1}$ is an automorphism because $\lambda_{p,q}$ does not belong to the spectrum of this restriction of A. Hence there exists unique $u' \in D^A_{p-1,q-1}$ such that $\lambda_{p,q} u' - \overline{A}u' = u$. By (16) this vector u' satisfies the equality (17) because of (14). Observe that the following correspondences described in this proof:

$$\begin{array}{c} \text{MHS} \quad \longmapsto \quad \text{strongly pseudo} - \text{real}\ \ \sigma - \text{operator} \\ \text{strongly pseudo} - \text{real}\ \ \sigma - \text{operator} \quad \longmapsto \quad \text{MHS} \end{array}$$

are inverse one of the other. □

References

- [1] G. Banaszak, J. Milewski, Hodge structures and Weierstrass sigma-function Comptes Rendus Acad. Sci. Paris, Ser. **I 350** (2012) 777-780
- [2] E. Cattani, A. Kaplan, W. Schmid, Degeneration of Hodge Structures, *Annals of Mathematics*, Second Series, **123**, No. 3 (1986), 457-535
- [3] P. Deligne, Structures de Hodge mixtes réeles, *Proceedings of Symposia in Pure Mathematics, American Mathematical Society* **55** (1994) 509-514
- [4] P. Deligne, On the SL(2)-orbits in Hodge theory, *IHES* pre-pub./M/82/58. (October 1982); the appendix ”Structures de Hodge mixtes réelles”, the Appendix to
- [5] B. Gordon, A survey of the Hodge conjecture for abelian varieties, Appendix B in “A survey of the Hodge conjecture”, by J. Lewis, 297–356 (1999)
- [6] C. Peters, J. Steenbrink, Mixed Hodge structures *Ergebnisse der Math. Springer* **52**, (2008).

Department of Mathematics and Computer Science, Adam Mickiewicz University, Poznań 61-614, Poland
E-mail address: `banaszak@amu.edu.pl`

Institute of Mathematics, Poznań University of Technology, ul. Piotrowo 3A, 60-965 Poznań, Poland
E-mail address: `jsmilew@wp.pl`